

REVIEW ARTICLE

# Not a one-trick pony: Diverse connectivity and functions of the rodent lateral geniculate complex

ABOOZAR MONAVARFESHANI,<sup>1,2</sup> UBADAH SABBAGH,<sup>1,3</sup> AND MICHAEL A. FOX<sup>1,2</sup>

<sup>1</sup>Developmental and Translational Neurobiology Center, Virginia Tech Carilion Research Institute, Roanoke, Virginia

<sup>2</sup>Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia

<sup>3</sup>Graduate Program in Translational Biology, Medicine, and Health, Virginia Tech, Blacksburg, Virginia

(RECEIVED January 20, 2017; ACCEPTED May 11, 2017)

## Abstract

Often mislabeled as a simple relay of sensory information, the thalamus is a complicated structure with diverse functions. This diversity is exemplified by roles visual thalamus plays in processing and transmitting light-derived stimuli. Such light-derived signals are transmitted to the thalamus by retinal ganglion cells (RGCs), the sole projection neurons of the retina. Axons from RGCs innervate more than ten distinct nuclei within thalamus, including those of the lateral geniculate complex. Nuclei within the lateral geniculate complex of nocturnal rodents, which include the dorsal lateral geniculate nucleus (dLGN), ventral lateral geniculate nucleus (vLGN), and intergeniculate leaflet (IGL), are each densely innervated by retinal projections, yet, exhibit distinct cytoarchitecture and connectivity. These features suggest that each nucleus within this complex plays a unique role in processing and transmitting light-derived signals. Here, we review the diverse cytoarchitecture and connectivity of these nuclei in nocturnal rodents, in an effort to highlight roles for dLGN in vision and for vLGN and IGL in visuomotor, vestibular, ocular, and circadian function.

**Keywords:** Thalamus, Retinal ganglion cells, Retinogeniculate, Lateral geniculate nucleus, Intergeniculate leaflet

## Introduction

Thalamus, from the Greek word *thalamos* meaning “inner chamber”, is a multifunctional diencephalic brain structure that plays important roles in receiving, processing, and relaying sensory information. The multiple and diverse functional roles of the thalamus may be best exemplified by those thalamic regions associated with light-derived visual stimuli. These regions receive, process, and relay not only classical image-forming visual information, which is a fundamental building block of vision, but also the less-well-studied nonimage-forming visual information.

Light-derived signals are detected and converted into neural signals by retinal photoreceptors. After being relayed and processed by interneurons in the inner nuclear layer (i.e., bipolar cells, horizontal cells, and amacrine cells), light-derived signals are transmitted to retinorecipient nuclei within the brain by retinal ganglion cells (RGCs). In both nocturnal and diurnal rodents, RGCs innervate approximately forty different retinorecipient regions, more than ten of which are located within the thalamus (Morin & Studholme, 2014; Martersteck et al., 2017) (Fig. 1). In nocturnal rodents, thalamic nuclei directly innervated by retinal axons include the dorsal lateral geniculate nucleus (dLGN),


ventral lateral geniculate nucleus (vLGN), intergeniculate leaflet (IGL), lateral posterior thalamic nucleus (LP; analogous to the pulvinar in higher mammals), anterodorsal thalamic nucleus (AD), centrolateral thalamic nucleus (CL), para-habenular zone (PHb), peripeduncular nucleus (PP), zona incerta (ZI), and subgeniculate nucleus (SubG) (Morin & Studholme, 2014) (Fig. 1). In addition to these retinorecipient regions, some thalamic nuclei, such as the thalamic reticular nucleus (TRN), process visual information but do not receive direct input from RGCs. Despite the plethora of retinorecipient targets in rodent thalamus, the vast majority of retinorecipient axons innervate one of three adjacent thalamic nuclei in the lateral geniculate complex [dLGN, IGL, and vLGN (Fig. 2)], each of which exhibits unique cytoarchitecture, circuitry, and function. Here, we review the unique properties of the lateral geniculate complex nuclei in nocturnal rodents, demonstrating the diverse roles thalamic nuclei exert in processing and transmitting light-derived information.

## Cytoarchitectural organization of retinorecipient thalamic nuclei

### dLGN

The dLGN receives, processes, and relays classical image-forming visual information and, for this reason, has received the most attention of all retinorecipient thalamic nuclei. In higher mammals, the

Address correspondence to: Michael A. Fox, Virginia Tech Carilion Research Institute, Director, Developmental and Translational Neurobiology Center, VTCRI, 2 Riverside Circle, Roanoke, VA 24016. E-mail: mafox1@vtc.vt.edu




**Fig. 1.** Retinorecipient nuclei of nocturnal rodents. This schematic illustrates the variety and distribution of brain nuclei innervated by retinal ganglion cells (see Morin and Studholme, 2014). Thalamic retinorecipient nuclei are colored in orange; other retinorecipient regions are colored gray. dLGN, dorsal lateral geniculate nucleus; IGL, intergeniculate leaflet; vLGNe, ventral lateral geniculate nucleus, external division; AD, anterodorsal thalamic nucleus; LP, lateral posterior thalamic nucleus; CL, CL; PP, PP; PHb, para-habenular zone; ZI, zona incerta; SubG, subgeniculate nucleus; SGN, suprageniculata nucleus; SON, supraoptic nucleus; SCN, suprachiasmatic nucleus; RCH, retrochiasmatic area; SBPV, subparaventricular zone; AHN, anterior hypothalamic area; LHA, lateral hypothalamic area; MeA, medial amygdala, anterior; MePV, medial amygdala, posteroventral; AAV, anterior amygdaloid area, ventral; SI, substantia innominata; MT, medial terminal nucleus; LT, lateral terminal nucleus; DT, dorsal terminal nucleus; PN, paranigral nucleus; MRN, midbrain reticular nucleus; PB, parabrachial nucleus; DR, dorsal raphe nucleus; PAG, periaqueductal gray; CPT, commissural pretectal nucleus; MPT, medial pretectal nucleus; PPT, posterior pretectal nucleus; APT, anterior pretectal nucleus; OPN, olivary pretectal nucleus; NOT, nucleus of OT; SC, superior colliculus; DCIC, dorsal cortex of the inferior colliculus; RGC, retinal ganglion cell.

dLGN has a distinctive cytoarchitecture with layers that receive eye- and function-specific retinal inputs. While cells in the dLGN of highly visual diurnal rodents, such as squirrels, are separated into at least five layers (Kaas et al., 1972; Van Hooser & Nelson, 2006), the dLGN of nocturnal rodents lacks gross cytoarchitecture lamination, despite having eye-specific domains (Reese & Cowey, 1983; Godement et al., 1984; Muir-Robinson et al., 2002; Jaubert-Miazza et al., 2005) (Fig. 2A). Nevertheless, evidence that the dLGN of rats (Reese, 1988), mice (Grubb & Thompson, 2004; Krahe et al., 2011), and hamsters (Emerson et al., 1982) are not anatomically homogenous emerged. The possibility that “hidden laminae” existed in rodent dLGN first arose from studies demonstrating the dorsolateral “shell” and ventromedial “core” regions of rodent dLGN contain populations of retinal terminals that are morphologically separable (Erzurumlu et al., 1988; Reese, 1988; Hammer et al., 2015). Hidden laminae have become more apparent with techniques that label individual classes of RGCs, of which there are more than thirty (Sanes & Masland, 2015; Baden et al., 2016). A series of studies in transgenic reporter mice (each of which labels a single class of RGCs) have demonstrated the existence of several RGC class-specific retinorecipient sublaminae in dLGN (Kim et al., 2008; Huberman et al., 2008a; Huberman et al., 2009; Kim et al., 2010; Hong & Chen, 2011; Kay et al., 2011) (Fig. 2D).

In addition to a heterogeneous distribution of retinal afferents, neuronal subtypes within the rodent dLGN are differentially distributed. Two main types of neurons exist in dLGN, both of which

are innervated by retinal afferents. Principal neurons, or thalamocortical (TC) relay cells, are excitatory projection neurons that originate from the caudal progenitor domain within the thalamic ventricular zone (i.e., prosomer 2) (Altman & Bayer, 1989; Puelles & Rubenstein, 2003; Vue et al., 2007) during rodent embryogenesis. Ultimately TC progenitor cells differentiate into at least three morphologically distinct classes in nocturnal rodents—biconical X-like cells, symmetrical Y-like cells, and hemispheric W-like TC cells (Krahe et al., 2011; Ling et al., 2012). These classes of TC relay cells closely resemble those reported in cats (Friedlander et al., 1981) and higher mammals (Irvin et al., 1993). Just as classes of relay cells are differentially distributed in cat and primate dLGN (Sherman, 1985; Nassi & Callaway, 2009), they are uniquely distributed in mouse dLGN: W-like cells occupy the dorsolateral shell of mouse dLGN and X- and Y-like cells occupy the ventromedial dLGN core (Krahe et al., 2011). While all three classes of TC relay cells project axons to visual cortex, recent evidence has demonstrated that regionally restricted cell types participate in functionally distinct parallel visual pathways in mice (Cruz-Martín et al., 2014; Bickford et al., 2015).

In addition to principal relay cells, rodent dLGN contains a small percentage (10–20%) of inhibitory interneurons, a cell type absent from most other dorsal thalamic regions (Arcelli et al., 1997; Jaubert-Miazza et al., 2005). The arrival of these interneurons occurs postnatally, after retinal inputs have targeted dLGN, formed immature connections, and begun to undergo activity-dependent refinement



**Fig. 2.** Organization of retinal projections in nuclei of the lateral geniculate complex. (A) Coronal view of a Nissl-stained mouse brain. Arrows indicate the location of dLGN, IGL, and vLGN. Image is from the Allen Brain Atlas (<http://www.brain-map.org>). (B–D) Schematic representation of coronal section through the lateral geniculate complex of nocturnal rodents. (B) depicts eyespecific segregation of retinal projections in dLGN, IGL, and vLGN. Terminals of ipsilateral retinal projections are depicted as green dots; terminals of contralateral retinal projections are depicted as orange dots. RGCs from which these projections arise are shown in the retinal cross sections. Dotted line in dLGN depicts the approximate boundary separating the dorsolateral shell (s) from the internal layer (i). The dotted line in vLGN depicts the boundary separating the external layer (e) from the internal layer (i). (C) depicts topographic mapping of retinal arbors in dLGN, vLGN, and IGL. Colors represent temporal (T) to nasal (N) location of RGCs in the retina (Feldheim et al., 1998; Pfeiffenberger et al., 2006; Huberman et al., 2008a). (D) depicts class-specific targeting of RGC axons to distinct sublaminae of dLGN, vLGN, and IGL. Colors represent some classes of RGCs studied with transgenic reporter mice. Names of these reporter mouse lines are indicated in parentheses (see Hattar et al. 2006; Kim et al. 2008; Huberman et al. 2009; Kim et al. 2010; Osterhout et al. 2011; Rivlin-Etzion et al. 2011). Color-filled dots in the dLGN, IGL, and vLGN represent retinal terminals (and are not meant to indicate that these terminals innervate distinct cells). dLGN, dorsal lateral geniculate nucleus; IGL, intergeniculate leaflet; vLGN, ventral lateral geniculate nucleus; dsRGC, direction-selective retinal ganglion cell; ipRGC, intrinsically photosensitive retinal ganglion cell.

(Jones & Rubenstein, 2004; Singh et al., 2012; Golding et al., 2014; Jager et al., 2016). The precise origin of these interneurons is currently under debate, with studies suggesting they arise from a rostral progenitor domain within the thalamus (i.e., prosomer 3) or from tectum (Virolainen et al., 2012; Golding et al., 2014; Jager et al., 2016). Evidence is also emerging that dLGN interneurons are not a homogeneous population in mice, and can instead be divided into at least two classes based on soma size, membrane capacitance, and neuronal nitric oxide synthase (nNOS) expression (Leist et al., 2016). Similar interneuron diversity has been reported in rats (Gabbott & Bacon, 1994b), cats (Montera & Zempel, 1985; Montero & Singer, 1985), and primates (Braak & Bachmann, 1985). At present, however, it remains unclear whether classes of local inhibitory interneurons exhibit regional preferences in the nocturnal rodent dLGN.

#### *vLGN and IGL*

Unlike dLGN, the vLGN of nocturnal rodents is organized into at least two easily identifiable laminae—the magnocellular external vLGN (vLGNe), which contains large cells and receives dense innervation from retina, and the parvocellular internal vLGN (vLGNi), which receives little, if any, retinal input (Niimi et al., 1963; Hickey & Spear, 1976; Gabbott & Bacon, 1994a; Harrington, 1997). These laminae are separated by a small neuron-free, fiber-rich neuropil. Cell types within vLGN are vastly different than those in dLGN resulting in a significant difference in the transcriptome of each region (Su et al., 2011; Yuge et al., 2011). Moreover, the vLGN lacks stereotypic TC relay cells, and has only a limited number of vesicular glutamate transporter-expressing glutamatergic neurons (Freneau et al., 2001; Yuge et al., 2011). Thus, in contrast to dLGN where glutamatergic neurons are the major cell type, vLGN contains a vastly higher population of GABAergic neurons (Gabbott & Bacon, 1994b; Harrington, 1997; Inamura et al., 2011). These major cellular differences reflect distinct embryonic origins of cells in vLGN, which are derived from progenitors in the most caudal prethalamus, a rostral domain of the thalamic ventricular zone, and the zona limitans interthalamica (ZLI) (Vue et al., 2007; Delaunay et al., 2009; Nakagawa & Shimogori, 2012; Virolainen et al., 2012). Although detailed studies of vLGN neurons lag behind similar characterizations of glutamatergic TC relay cells and local GABAergic interneurons in dLGN, expression studies and transgenic reporter mice strongly suggest that distinct classes of cells are distributed in a laminar arrangement in vLGNe, even if it is not apparent at the gross cytoarchitectural level (Moore & Card, 1994; <http://www.brain-map.org>).

Although vLGN and IGL likely serve different functions in nocturnal rodents and are anatomically distinguishable (Hickey & Spear, 1976; Moore & Card, 1994; Morin, 2013), we have grouped them together throughout this review because of shared features that will be discussed. Like those in vLGNe, neurons in the IGL originate from a rostral region of the thalamic progenitor zone and the ZLI (Vue et al., 2007; Delaunay et al., 2009). A large fraction of neurons in IGL generate GABA (Moore & Speh, 1993), few are glutamatergic, and none project axons to visual cortex (Harrington, 1997). However, it is important to highlight that some classes of IGL neurons are absent from vLGN. This includes NPY-expressing neurons in rodent IGL that project axons to hypothalamic nuclei (Card & Moore, 1989; Moore & Card, 1994; Harrington, 1997). Based on available data, classes of morphologically and neurochemically distinct neurons do not appear to cluster into distinct layers or regions of IGL, except, perhaps, for coarse differences in

their distribution between rostral and caudal regions of the IGL (Brauer et al., 1983; Moore & Card, 1994; Morin, 2013).

#### **Afferent projections of retinorecipient thalamic nuclei**

##### *dLGN: Retinal afferents*

In mammals, the primary excitatory drive onto TC relay cells is provided by retinal inputs (Sherman, 2005; Petrof & Sherman, 2013). Anatomically, retinal projections to dLGN are spatially organized in (at least) three fundamental ways. First, retinal afferents are segregated into nonoverlapping eye-specific domains in an activity-dependent manner (Huberman et al., 2008a; Zhang et al., 2012). The dLGN of nocturnal rodents receives a relatively small contribution (5–10%) of retinal afferents from the ipsilateral retina and these projections are confined to a ventromedial core region of the dLGN (Jaubert-Miazza et al., 2005; Gaillard et al., 2013; Morin & Studholme, 2014) (Fig. 2B). Second, retinal projections to dLGN are organized topographically—so that neighboring RGCs provide input to neighboring TC relay cells and provide a continuous and faithful representation of spatial information from retina to brain (Feldheim et al., 1998; Pfeiffenberger et al., 2006; Huberman et al., 2008a; Cang & Feldheim, 2013) (Fig. 2C). Third, and perhaps most remarkably, retinal projections undergo class-specific segregation in rodent dLGN (Fig. 2D). Although more than 30 classes of RGCs exist in nocturnal rodents only a subset innervate dLGN (Sanes & Masland, 2015; Baden et al., 2016; Ellis et al., 2016). This suggests that targeting mechanisms exist that guide some classes of RGC axons into dLGN and exclude others. Once appropriate classes of retinal axons enter dLGN, they are further segregated into a newly-appreciated laminar organization (Hong & Chen, 2011; Dhande & Huberman, 2014; Sanes & Masland, 2015). The presence of these stereotyped class-specific retinal projections has been elegantly revealed by transgenic reporter mice in which individual RGC classes are labeled with reporter proteins (Hattar et al., 2006; Kim et al., 2008; Huberman et al., 2008b; Huberman et al., 2009; Kim et al., 2010; Kay et al., 2011; Rivlin-Etzion et al., 2011; Dhande et al., 2013; Triplett et al., 2014). While only a small set of individual RGC projections have been mapped with this approach, some rules are beginning to emerge. First, projections of direction-selective classes of RGCs arborize in more dorsolateral regions of dLGN, including the shell of dLGN (Kim et al., 2008; Huberman et al., 2009; Rivlin-Etzion et al., 2011; Cruz-Martín et al., 2014). Second, there is considerable overlap in the laminar termination zones of RGC axons (Fig. 2D), and taken in the context of recent ultrastructural and circuit tracing experiments in dLGN (Morgan et al., 2016; Rompani et al., 2017), this raises the possibility that individual TC relay cells may receive inputs from multiple classes of RGCs.

In addition to being segregated based on eye of origin, topography, and RGC class, retinal inputs in dLGN are structurally and functionally distinct from retinal inputs in other retinorecipient nuclei, even other thalamic nuclei (Sherman, 2005; Hammer et al., 2014). Specifically, retinal terminals onto dLGN TC relay cells are significantly larger than all other terminals in dLGN (and larger than retinal terminals in all other retinorecipient nuclei), and exhibit unique ultrastructural morphology and function (Guillery, 1969; Lund & Cunningham, 1972; Sherman, 2004; Guido, 2008; Bickford et al., 2010; Hong & Chen, 2011). It is worth pointing out, however, that at least two distinct types of RG synapses have been identified in rodent dLGN: “simple encapsulated” RG synapses, in which a single

retinal terminal synapses onto a TC relay cell dendrite, and “complex encapsulated” RG synapses in which axons from numerous RGCs converge to innervate adjacent regions of a TC relay cell dendrite (Lund & Cunningham, 1972; Hammer et al., 2015; Morgan et al., 2016). Finally, it is important to point out that retinal projections not only innervate TC relay cells, but also local interneurons in nocturnal rodents (Sherman, 2004; Seabrook et al., 2013b).

#### *dLGN: Nonretinal afferents*

While retinal inputs provide the excitatory drive to TC relay cells, they account for only 5–10% of the total inputs onto a relay cell and are far outnumbered by nonretinal inputs (Sherman & Guillery, 2002; Bickford et al., 2010; Cetin & Callaway, 2014). A summary of the main inputs to rodent dLGN is depicted in Fig. 3.

While many nonretinal inputs onto dLGN TC relay cells have modulatory or inhibitory roles, a recent study identified a novel glutamatergic nonretinal source of “driver-like” input onto dLGN TC relay cells (Bickford et al., 2015). These inputs arise from the ipsilateral superior colliculus (SC) and terminate onto W-like TC relay cells in the dorsolateral shell of dLGN (Harting et al., 1991a; Bickford et al., 2015). Circuit tracing experiments indicate these excitatory tectogeniulate connections contribute to the processing and transmission of direction-selective visual information (Bickford et al., 2015).

While tectogeniulate inputs represent a minor source of inputs to dLGN, a more significant portion of nonretinal glutamatergic inputs arise from cortical projection neurons in layer VI of primary visual cortex (Sherman, 2016). Corticothalamic inputs are small, located on distal portions of TC relay cell dendrites, and generate weak excitatory postsynaptic potentials (EPSPs) in relay cells (Sherman & Guillery, 2002; Petrof & Sherman, 2013). For this reason, it is likely that these inputs are insufficient for the relay of information alone and are, therefore, modulatory in nature (Petrof & Sherman, 2013). Despite these features, corticothalamic inputs do significantly influence RG transmission by affecting the gain of signal transmission and sharpening of receptive field properties of TC relay cells (Sherman & Guillery, 2002; Briggs & Usrey, 2008; Olsen et al., 2012; Bickford, 2015).

In addition to tectal and cortical glutamatergic inputs, TC relay cells in higher mammals receive modulatory cholinergic, serotonergic, noradrenergic, and dopaminergic inputs from a variety of sources in the brainstem including parabrachial nucleus, pedunculopontine region, locus coeruleus, dorsal raphe nucleus of the midbrain, and the midbrain reticular formation (Mackay-Sim et al., 1983; de Lima et al., 1985; De Lima & Singer, 1987; Papadopoulos & Parnavelas, 1990a, b; McCormick, 1992; Jones, 2012). At present, some of these afferent projections have been demonstrated in nocturnal rodents (Hallanger et al., 1987; Harting et al., 1991b), but additional studies are needed to map specific sources of these inputs and to understand their role in signal processing in rodents.

Finally, the last significant source of nonretinal inputs to dLGN are inhibitory GABAergic inputs that arise from both local inhibitory neurons and projection neurons in the TRN, a region that forms a lateral shell around dorsal thalamus in nocturnal rodents (Hale et al., 1982; Guillery & Harting, 2003; Pinaut, 2004). Inhibitory neurons in the ipsilateral pretectum also project to dLGN (Born & Schmidt, 2007), however evidence suggests that these projections innervate dLGN interneurons not TC relay cells (Wang et al., 2002; Born & Schmidt, 2007), adding further complexity to dLGN circuitry.

An interesting facet of the convergence of retinal and nonretinal inputs in dLGN is that their development appears tightly coordinated.


Retinal axons target and innervate dLGN prior to the arrival of non-retinal inputs and play instructive roles in the establishment of nonretinal circuitry (Brooks et al., 2013; Seabrook et al., 2013a; Golding et al., 2014; Grant et al., 2016). Likewise, nonretinal inputs contribute to the development and function of retinogeniulate synapses. For example, the presence of corticothalamic axons and corticogeniulate synapses play essential roles in the establishment, refinement, and maintenance of retinal inputs (Shanks et al., 2016; Thompson et al., 2016).

#### *vLGN and IGL: Retinal afferents*

Several features of retinal projections to vLGN are similar to those in dLGN: retinal afferents provide a main excitatory drive to vLGN; the arrival of retinal inputs in vLGN occurs neonatally and precedes nonretinal inputs (Su et al., 2011); retinal inputs are mapped topographically in vLGN and these inputs are segregated into nonoverlapping eye-specific domains (Holcombe & Guillery, 1984; Hammer et al., 2014; Morin & Studholme, 2014). On this last point, it warrants mention that ipsilateral retinal projections occupy a region of vLGN that is more complex and less stereotyped than its counterparts in dLGN. For this reason, activity-dependent refinement of eye-specific retinal projections has not been thoroughly characterized in vLGN, nor has it served as an anatomical readout of activity-dependent refinement as has been the case in dLGN (Jaubert-Miazza et al., 2005; Demas et al., 2006; Stevens et al., 2007; Huberman et al., 2008b; Rebsam et al., 2009; Xu et al., 2011; Rebsam et al., 2012; Dilger et al., 2015).

There are, however, several dramatic differences between retinal projections in dLGN and vLGN. First, the majority of dLGN-projecting classes of RGCs fail to send collateral axon branches into vLGN despite having to pass by it (or through it) on the way to dLGN (Huberman et al., 2008b; Kim et al., 2008; Huberman et al., 2009; but see also; Rivlin-Etzion et al., 2011). Instead, sets of nonimage-forming classes of RGCs, including the M1 class of melanopsin-expressing intrinsically photosensitive RGCs (ipRGCs; labeled in *Opn4<sup>taulacz/taulacz</sup>* mice) and RGCs labeled in *Cdh3-GFP* mice, innervate vLGN (Hattar et al., 2006; Osterhout et al., 2011). It is worth noting that a small subset of ipRGC terminals have been reported in the medial-most region of dLGN in *Opn4<sup>taulacz/taulacz</sup>* mice. While these may reflect projections from M1 ipRGCs, it is also possible that they belong to axons from other ipRGCs, such as M3 ipRGCs (Schmidt et al., 2011). Indeed other classes of ipRGCs innervate dLGN and mediate image-forming visual signals in nocturnal rodents (Ecker et al., 2010; Estevez et al., 2012). Projections of M1 ipRGCs and *Cdh3-GFP* RGCs arborize broadly across all regions of vLGN, raising questions as to whether hidden lamina exist in vLGN. However, it is important to point out that studies in diurnal rats identified distinct lamination of retinal projections in this region (Gaillard et al., 2013). Moreover, only a small set of the RGCs that likely innervate rodent vLGN have been identified and studied to date, leaving open the possibility that additional RGC classes will be identified whose projections are regionally restricted to specific sublamina of vLGN. In support of this, there is a small region of vLGN underlying the optic tract (OT) that contains a region with morphologically distinct retinal arbors (Hammer et al., 2014), much like that observed in the dorsolateral shell of dLGN (Reese, 1988; Grubb & Thompson, 2004).

Significant differences also exist in the anatomy and physiology of retinal synapses in vLGN compared to dLGN. Retinal terminals in rodent vLGN are remarkably smaller and less morphologically



**Fig. 3.** Afferent and efferent projections of rodent dLGN. **(A)** Sources of afferent projections to dLGN are colored green. **(B)** Brain regions innervated by dLGN efferents are colored in orange. dLGN, dorsal lateral geniculate nucleus; vLGN, ventral lateral geniculate nucleus; TRN, thalamic reticular nucleus; SC, superior colliculus; PRT, pretectal region; MRN, midbrain reticular nucleus; DR, dorsal raphe nucleus; PPN, pedunculopontine nucleus; PBG, parabrachial nucleus; PB, parabrachial nucleus; LC, locus coeruleus; VIS1, visual cortex, layer I; VIS4, visual cortex, layer IV; VIS6, visual cortex, layer VI; RGC, retinal ganglion cell.

complex than those in dLGN, and this difference is reflected in their ability to elicit considerably weaker EPSPs (Mize & Homer, 1984; Hammer et al., 2014). While several features of retinogeniculate synapses in vLGNs more closely resemble features of modulatory glutamatergic inputs (including their size, synaptic strength, and relative level of convergence on postsynaptic neurons), retinal inputs onto vLGN principal neurons exhibit paired-pulse depression and are likely to be “driver” inputs (Hammer et al., 2014). Ultrastructural analysis further suggests that retinal terminals in vLGNs do not form “complex encapsulated” RG synapses and are

not typically ensheathed by glial processes (Stelzner et al., 1976; Hammer et al., 2014), both of which are features of RG synapses in rodent dLGN (Hammer et al., 2014; Hammer et al., 2015). These differences in retinal input type, synaptic morphology, and synaptic physiology suggest that light-derived information is processed differently in vLGNs.

Characteristics of retinal afferents in the nocturnal rodent IGL diverge even farther from their analogues in dLGN. Retinal projections to IGL are not segregated into nonoverlapping eye-specific domains (Su et al., 2013; Hammer et al., 2014; Morin & Studholme, 2014)

nor are they topographically mapped (Harrington, 1997) (Fig. 2). These two features of retinal inputs distinguish IGL from vLGNe, which is surprising given that the only known classes of RGCs that project to IGL, M1 ipRGCs, and Cdh6-expressing RGCs, also project to vLGNe (Hattar et al., 2006; Osterhout et al., 2011). It is possible that subsets of RGCs in these classes differentially target IGL and vLGNe, or that each retinorecipient region has unique target-derived signals that allow for differential nonimage-forming RGC axon targeting mechanisms (Fox & Guido, 2011).

Retinal terminals in IGL do share ultrastructural similarities with retinal terminals in dLGN and vLGNe, such as the presence of pale mitochondria and round synaptic vesicles (Moore & Card, 1994). In contrast to retinal terminals in dLGN, which reside on large proximal dendrites (Rafols & Valverde, 1973; Wilson et al., 1984), retinal inputs appear to contact small diameter, distal dendrites in IGL (Moore & Card, 1994). Analysis of retinal terminal size by anterograde labeling suggests that these terminals are similar in size to those in vLGNe, but are much less densely distributed than those in other major retinorecipient nuclei (Hammer et al., 2014). Some peculiar features of retinal terminals in IGL have been noted in the literature and warrant mention here. For example, while retinal terminals in vLGNe, dLGN, and all other retinorecipient nuclei contain Vesicular Glutamate Transporter 2 (VGluT2), little, if any, of this transporter (or its closely related family member VGluT1) is present in IGL (Fujiyama et al., 2003; Su et al., 2013; Hammer et al., 2014). The limited level of VGluT expression in IGL appears unchanged following surgical or genetic enucleation (Fujiyama et al., 2003; Hammer et al., 2014), further suggesting retinal terminals in IGL may lack machinery for glutamate release. Despite this apparent lack of machinery for glutamate packaging and release, Blasiak et al. (2009) showed that glutamate receptor antagonists impair excitatory responses to OT stimulation in IGL neurons. How do we reconcile such differences? It is possible that VGluT2 is present in retinal terminals in IGL but is below the limit of detection by immunostaining. Would such low levels of vGluT2 allow faithful transmission of signals at retinal synapses in IGL? Perhaps. Retinal terminals in IGL contain pituitary adenylate cyclase-activating peptide (PACAP) (Engelund et al., 2010), a neurotransmitter that can be co-released with glutamate and amplifies glutamatergic signaling (Kopp et al., 2001; Michel et al., 2006). Co-release of PACAP may reduce the necessity of retinal terminals to package and release large quantities of glutamate and therefore retinal terminals may require minimal levels of VGluTs.

In addition to differences in synapse size and neurotransmitters released, a final asymmetry between retinal connections in IGL and other retinorecipient nuclei is that a considerable fraction of principle projection neurons in IGL (such as NPY + neurons) are not directly innervated by retinal inputs. Instead, the ability of these projection neurons to propagate light-derived signals requires yet-to-be identified interneurons (Thankachan & Rusak, 2005; Morin, 2013), and suggests the existence of a unique circuitry for the transmission of sensory information in IGL.

#### *vLGN and IGL: Nonretinal afferents*

Despite some differences in retinal connectivity with vLGNe and IGL, there appears to be a high degree of similarity in the nonretinal afferents innervating these structures. It is worth noting three distinctions between sources of nonretinal afferents to vLGNe and IGL when compared with dLGN. First, the number of nuclei that project afferents to rodent vLGNe and IGL far exceed and are far

more diverse than those to dLGN (Figs. 4 and 5). An incomplete list of these sources includes superior colliculus, visual cortex, olivary pretectal nucleus, anterior pretectal nucleus, posterior pretectal nucleus, locus coeruleus, dorsal raphe nucleus, subparafascicular thalamic nucleus, mesencephalic nucleus, lateral dorsal tegmental nucleus, medial and lateral terminal nuclei, supraoculomotor periaqueductal gray, retrorubral nucleus, pontine reticular nucleus, parabrachial nucleus, and medial vestibular nucleus (Cosenza & Moore, 1984; Moore & Card, 1994; Moore et al., 2000; Vrang et al., 2003; Horowitz et al., 2004) (Figs. 4 and 5). In addition, neurons in rodent IGL (but not vLGNe) receive afferents from a variety of sources which include (but are not limited to) prefrontal cortex, ZI, suprachiasmatic nucleus, cuneiform nucleus, and superior and lateral vestibular nuclei (Morin & Blanchard, 1998, 1999; Vrang et al., 2003; Horowitz et al., 2004) (Fig. 5). Second, subcortical projections to vLGNe and IGL arise from both ipsilateral and contralateral sources in nocturnal rodents (Moore & Card, 1994; Vrang et al., 2003; Horowitz et al., 2004). For example, IGL neurons receive bilateral input from the olivary pretectal nucleus (OPN) and suprachiasmatic nucleus (SCN), and contralateral inputs from the other IGL (Vrang et al., 2003); likewise, vLGNe neurons receive input from the contralateral vLGNe (Cosenza & Moore, 1984). Third, few sources of afferents are shared with dLGN, strongly inferring diverse roles of these thalamic regions in processing light-derived information (Horowitz et al., 2004). Even in cases when projections to all three regions of the lateral geniculate complex arise from a single brain region, they originate from different cellular sources. For example while dLGN, vLGNe, and IGL all receive input from corticothalamic cells in primary visual cortex, inputs to dLGN arise from layer VI cells whereas those innervating vLGNe and IGL arise from layer V (Cosenza & Moore, 1984; Bourassa & Deschênes, 1995; Jacobs et al., 2007; Seabrook et al., 2013a; Hammer et al., 2014).


### **Efferent projections of retinorecipient thalamic nuclei**

#### *dLGN efferents*

Of the retinorecipient thalamic nuclei discussed here, the dLGN has by far the fewest targets of efferent projections. In fact, the relative simplicity of efferent projections from dLGN is striking and emphasizes a singular function of dLGN in processing and transferring image-forming visual information. TC relay cells project axons to only two ipsilateral regions in nocturnal rodents: primary visual cortex and TRN (Rafols & Valverde, 1973; Towns et al., 1982; Reese & Cowey, 1983; Crabtree & Killackey, 1989; López-Bendito & Molnár, 2003; Jurgens et al., 2012). Recent studies in mice have demonstrated that projections to visual cortex exhibit class-specificity, with W-like relay cells in the dorsolateral shell of dLGN, conveying direction-selective visual information to layer I and Y-like and X-like relay cells projecting to layer IV of primary visual cortex (Cruz-Martín et al., 2014; Bickford et al., 2015). At present, it remains unclear whether these three classes of TC relay cells make unique connections with TRN neurons in nocturnal rodents. It is worth mentioning that, in cat, Y-cells provide the predominant dLGN input to the perigeniculate nucleus, the visual sector of cat TRN which overlies dLGN (Dubin & Cleland, 1977; Friedlander et al., 1980).

#### *vLGN and IGL efferents*

Efferents from vLGNe and IGL target many of the same brain regions and, in contrast to the simplicity of the dLGN efferents,




**Fig. 4.** Afferent and efferent projections of rodent vLGN. **(A)** Sources of afferent projections to vLGN are depicted in green. **(B)** Brain regions innervated by vLGN neurons are depicted in orange. dLGN, dorsal lateral geniculate nucleus; vLGN, ventral lateral geniculate nucleus, external division; LP, lateral posterior thalamic nucleus; LD, lateral dorsal nucleus; MD, medial dorsal nucleus; ZI, zona incerta; RH, rhomboid nucleus; RE, reuniens nucleus; SMT, submedial nucleus of the thalamus; LHA, lateral hypothalamic area; PH, posterior hypothalamic nucleus; SC, superior colliculus; PRT, pretectal region; MRN, midbrain reticular nucleus; PAG, periaqueductal gray; DR, dorsal raphe nucleus; PPN, pedunculopontine nucleus; RR, retrorubral nucleus; MT, medial terminal nucleus; LT, lateral terminal nucleus; DT, dorsal terminal nucleus; LC, locus coeruleus; PB, parabrachial nucleus; Bar, Barrington's nucleus; MV, medial vestibular nucleus; LDT, laterodorsal tegmental nucleus; PRNc, pontine reticular nucleus; MDRN, medullary reticular nucleus; IO, accessory inferior olivary nucleus; VIS5, visual cortex, layer 5; RGC, retinal ganglion cell.

are diverse and far-reaching. Importantly, neither vLGN nor IGL project efferents to visual cortex or any other cortical region (Harrington, 1997). Instead, their efferents innervate regions that regulate visuomotor function, eye movement, vestibular function, and circadian function (Moore et al., 2000) (Figs. 4 and 5). A noteworthy and major target of efferents from vLGN and IGL is the superior colliculus, a region intimately involved in multisensory

integration, visuomotor function, and coordination of eye movements. Projections from vLGN and IGL arborize in all layers of the ipsilateral SC and represent the largest thalamic source of afferents to SC in nocturnal rodents (Matute & Streit, 1985; Taylor et al., 1986). However, whether these projections are excitatory, inhibitory, or modulatory remains unresolved. Efferents from both vLGN and/or IGL also target other midbrain structures,





**Fig. 5.** Afferent and efferent projections of rodent IGL. (A) Sources of afferent projections to IGL are colored green. (B) Brain regions innervated by IGL neurons are depicted in orange. IGL, intergeniculate leaflet; LP, lateral posterior thalamic nucleus; PP, PP; ZI, zona incerta; SPF, subparafascicular nucleus; RH, rhomboid nucleus; RE, reuniens nucleus; PVT, paraventricular nucleus; SCN, suprachiasmatic nucleus; RCH, retrochiasmatic area; SBPV, subparaventricular zone; AHN, anterior hypothalamic area; PH, posterior hypothalamic nucleus; VMH, ventromedial hypothalamic nucleus; DMH, dorsomedial nucleus of the hypothalamus; SC, superior colliculus; PRT, pretectal region; MT, medial terminal nucleus; LT, lateral terminal nucleus; DT, dorsal terminal nucleus; DR, dorsal raphe nucleus; PAG, periaqueductal gray; CUN, cuneiform nucleus; PPN, pedunculopontine nucleus; RR, retrorubral nucleus; LC, locus coeruleus; LDT, laterodorsal tegmental nucleus; Bar, Barrington's nucleus; MV, medial vestibular nucleus; LAV, lateral vestibular nucleus; SUV, superior vestibular nucleus; PRNc, pontine reticular nucleus; VIS5, visual cortex, layer 5; ACA5/6, anterior cingulate area, layer 5 and 6; RGC, retinal ganglion cell.

including the ipsilateral OPN, the nucleus of the OT, and the anterior pretectal nucleus (Cadusseau & Roger, 1991; Harrington, 1997; Moore et al., 2000). Other regions associated with eye movements, visuomotor function, and attention innervated by vLGN and IGL include two of the accessory optic system nuclei (lateral terminal nucleus and medial terminal nucleus; Swanson et al., 1974;

Ribak & Peters, 1975), zona incerta (Brauer & Schober, 1982), and nuclei within the pons (Harrington, 1997).

In addition to visuomotor functions, vLGN and IGL efferents project to both hypothalamic and thalamic regions (Moore et al., 2000). The largest of these projections innervates the SCN, and is referred to as the geniculohypothalamic (GH) tract. GH projections

are GABAergic, are important for modulating circadian function (Harrington, 1997), and originate from NPY<sup>+</sup> or Enk<sup>+</sup> neurons in the IGL (Card & Moore, 1982; Harrington et al., 1987; Card & Moore, 1989). A separate set of neurons in vLGN and IGL innervate contralateral vLGN and IGL (Harrington, 1997), contralateral dLGN (Mikkelsen, 1992; Kolmac et al., 2000), and, at least in higher mammals, pulvinar (Nakamura & Kawamura, 1988).

### Diverse connectivity leads to diverse functions of retinorecipient thalamic nuclei

Retinorecipient nuclei within the thalamus exemplify the diverse roles these brain regions play in sensory processing. Despite residing adjacent to each other in the lateral geniculate complex and receiving light-derived signals directly from the retina, there are few similarities in their cytoarchitecture, connectivity, or function of dLGN, vLGN, and IGL.

Based upon its efferent projections to cortex and early studies showing near unitary matching of retinal afferents to TC relay cells (Glees & le Gros Clark, 1941), the dLGN was initially characterized as a simple relay of visual information. Certainly, the relative simplicity of its efferent projections suggests a near singular role in transmitting image-forming visual information to visual cortex. However, describing the dLGN as a simple relay underestimates its role in processing image-forming visual information. Modulatory feedback from cortex, cholinergic inputs from a subset of brainstem nuclei, and inhibition from interneurons, endows the dLGN with the ability to shape visual information before transmitting it to higher cortical centers (Piscopo et al., 2013; Roth et al., 2016; Weyand, 2016; Rompani et al., 2017). For example, top-down feedback from corticothalamic inputs is thought to increase the selectivity of TC relay cells, sharpen TC relay cell receptive field properties, enhance synchronicity among TC relay cells, and influence the gain of retinogeniculate transmission (Sherman & Guillery, 2002; Briggs & Usrey, 2008; Weyand, 2016). In nocturnal rodents (and higher mammals) the manner in which retinal inputs innervate dLGN offers the possibility of modifying light-derived signals: feed forward inhibition of retinal inputs through local interneurons can either enhance temporal specificity of RG transmission or can enhance lateral inhibition (Martinez et al., 2014; Weyand, 2016), single retinal axons innervating multiple TC relay cells can amplify visual signals (Weyand, 2016), and the convergence of numerous retinal axons on single relay cell dendrites can produce TC receptive fields that are not present in retina (Hammer et al., 2015; Morgan et al., 2016; Weyand, 2016; Rompani et al., 2017). Thus, dLGN is more likely an active component of the machinery required to transform image-forming information into vision and not a passive relay.

In contrast, vLGN and IGL have little (if any) role in vision. Despite not innervating visual cortex, neurons in these thalamic regions do innervate visual system centers upstream of dLGN (e.g., SC) and even provide inputs to dLGN. Lesion experiments also suggest a potential role for vLGN and IGL in visual intensity discrimination (Horel, 1968; Legg & Cowey, 1977a, b; Harrington, 1997), however, these studies (and, in fact, all lesion studies of these regions) need to be interpreted cautiously as the lesions disrupt the overlying OT and may have secondary effects on other retinorecipient nuclei. Based upon input from classes of RGCs that convey nonimage-forming visual information, and projections to a variety of subcortical structures, roles for vLGN and IGL in modulating circadian rhythms, visuomotor function, eye movements, and vestibular function seem likely. While lesion studies have addressed

some of these possibilities in nocturnal rodents (Harrington & Rusak, 1986; Lewandowski & Usarek, 2002) more elegant and specific genetic approaches to lesion or silence activity in vLGN and IGL have yet-to-be applied. Such studies are needed to definitively identify the functions of vLGN and IGL, and to identify any potential differences in these two nuclei.

### Acknowledgments

We apologize to those whose work we excluded from this review due to space constraints. Work in the Fox laboratory is supported by the National Institutes of Health (EY021222, EY024712, AI124677) and by a Brain and Behavior Research Foundation NARSAD Independent Investigator Award. A.M. is supported by a Virginia Tech Carilion Research Institute Medical Research Scholar fellowship.

### References

- ALTMAN, J. & BAYER, S.A. (1989). Development of the rat thalamus: VI. The posterior lobule of the thalamic neuroepithelium and the time and site of origin and settling pattern of neurons of the lateral geniculate and lateral posterior nuclei. *Journal of Comparative Neurology* **284**, 581–601.
- ARCELLI, P., FRASSONI, C., REGONDI, M., BIASI, S. & SPREAFICO, R. (1997). GABAergic neurons in mammalian thalamus: A marker of thalamic complexity? *Brain Research Bulletin* **42**, 27–37.
- BADEN, T., BERENS, P., FRANKE, K., ROSÓN, M.R., BETHGE, M. & EULER, T. (2016). The functional diversity of retinal ganglion cells in the mouse. *Nature* **529**, 345–350.
- BICKFORD, M.E. (2015). Thalamic circuit diversity: Modulation of the driver/modulator framework. *Frontiers in Neural Circuits* **9**, 86.
- BICKFORD, M.E., SLUSARCZYK, A., DILGER, E.K., KRAHE, T.E., KUCUK, C. & GUIDO, W. (2010). Synaptic development of the mouse dorsal lateral geniculate nucleus. *The Journal of Comparative Neurology* **518**, 622–635.
- BICKFORD, M.E., ZHOU, N., KRAHE, T.E., GOVINDAIAH, G. & GUIDO, W. (2015). Retinal and tectal “driver-like” inputs converge in the shell of the mouse dorsal lateral geniculate nucleus. *The Journal of Neuroscience* **35**, 10523–10534.
- BLASIAK, A., BLASIAK, T. & LEWANDOWSKI, M. (2009). Electrophysiology and pharmacology of the optic input to the rat intergeniculate leaflet *in vitro*. *Acta Physiologica Polonica* **60**, 171.
- BORN, G. & SCHMIDT, M. (2007). GABAergic pathways in the rat subcortical visual system: A comparative study *in vivo* and *in vitro*. *European Journal of Neuroscience* **26**, 1183–1192.
- BOURASSA, J. & DESCHÊNES, M. (1995). Corticothalamic projections from the primary visual cortex in rats: A single fiber study using biocytin as an anterograde tracer. *Neuroscience* **66**, 253–263.
- BRAAK, H. & BACHMANN, A. (1985). The percentage of projection neurons and interneurons in the human lateral geniculate nucleus. *Human Neurobiology* **4**, 91–95.
- BRAUER, K. & SCHÖBER, W. (1982). Identification of geniculate-tectal relay neurons in the rat's ventral lateral geniculate nucleus. *Experimental Brain Research* **45**, 84–88.
- BRAUER, K., SCHÖBER, W., LEIBNITZ, L., WERNER, L., LÜTH, H. & WINKELMANN, E. (1983). The ventral lateral geniculate nucleus of the albino rat morphological and histochemical observations. *Journal für Hirnforschung* **25**, 205–236.
- BRIGGS, F. & USREY, W.M. (2008). Emerging views of corticothalamic function. *Current Opinion in Neurobiology* **18**, 403–407.
- BROOKS, J.M., SU, J., LEVY, C., WANG, J.S., SEABROOK, T.A., GUIDO, W. & FOX, M.A. (2013). A molecular mechanism regulating the timing of corticogeniculate innervation. *Cell Reports* **5**, 573–581.
- CADUSSEAU, J. & ROGER, M. (1991). Cortical and subcortical connections of the pars compacta of the anterior pretectal nucleus in the rat. *Neuroscience Research* **12**, 83–100.
- CANG, J. & FELDHEIM, D.A. (2013). Developmental mechanisms of topographic map formation and alignment. *Annual Review of Neuroscience* **36**, 51–77.
- CARD, J.P. & MOORE, R.Y. (1982). Ventral lateral geniculate nucleus efferents to the rat suprachiasmatic nucleus exhibit avian pancreatic polypeptide-like immunoreactivity. *Journal of Comparative Neurology* **206**, 390–396.

- CARD, J.P. & MOORE, R.Y. (1989). Organization of lateral geniculate-hypothalamic connections in the rat. *Journal of Comparative Neurology* **284**, 135–147.
- CETIN, A. & CALLAWAY, E.M. (2014). Optical control of retrogradely infected neurons using drug-regulated “TLoop” lentiviral vectors. *Journal of Neurophysiology* **111**, 2150–2159.
- COSENZA, R.M. & MOORE, R.Y. (1984). Afferent connections of the ventral lateral geniculate nucleus in the rat: An HRP study. *Brain Research* **310**, 367–370.
- CRABTREE, J.W. & KILLACKEY, H.P. (1989). The topographic organization and axis of projection within the visual sector of the rabbit's thalamic reticular nucleus. *European Journal of Neuroscience* **1**, 94–109.
- CRUZ-MARTÍN, A., EL-DANAF, R.N., OSAKADA, F., SRIRAM, B., DHANDE, O.S., NGUYEN, P.L., CALLAWAY, E.M., GHOSH, A. & HUBERMAN, A.D. (2014). A dedicated circuit links direction-selective retinal ganglion cells to the primary visual cortex. *Nature* **507**, 358–361.
- DE LIMA, A.D. & SINGER, W. (1987). The serotonergic fibers in the dorsal lateral geniculate nucleus of the cat: Distribution and synaptic connections demonstrated with immunocytochemistry. *Journal of Comparative Neurology* **258**, 339–351.
- DE LIMA, D.A., MONTERO, V. & SINGER, W. (1985). The cholinergic innervation of the visual thalamus: An EM immunocytochemical study. *Experimental Brain Research* **59**, 206–212.
- DELAUNAY, D., HEYDON, K., MIGUEZ, A., SCHWAB, M., NAVE, K.A., THOMAS, J.L., SPASSKY, N., MARTINEZ, S. & ZALC, B. (2009). Genetic tracing of subpopulation neurons in the prethalamus of mice (*Mus musculus*). *Journal of Comparative Neurology* **512**, 74–83.
- DEMAS, J., SAGDULLAEV, B.T., GREEN, E., JAUBERT-MIAZZA, L., MCCALL, M.A., GREGG, R.G., WONG, R.O. & GUIDO, W. (2006). Failure to maintain eye-specific segregation in nob, a mutant with abnormally patterned retinal activity. *Neuron* **50**, 247–259.
- DHANDE, O.S., ESTEVEZ, M.E., QUATTROCHI, L.E., EL-DANAF, R.N., NGUYEN, P.L., BERSON, D.M. & HUBERMAN, A.D. (2013). Genetic dissection of retinal inputs to brainstem nuclei controlling image stabilization. *The Journal of Neuroscience* **33**, 17797–17813.
- DHANDE, O.S. & HUBERMAN, A.D. (2014). Retinal ganglion cell maps in the brain: Implications for visual processing. *Current Opinion in Neurobiology* **24**, 133–142.
- DILGER, E.K., KRAHE, T.E., MORHARDT, D.R., SEABROOK, T.A., SHIN, H-S. & GUIDO, W. (2015). Absence of plateau potentials in dLGN cells leads to a breakdown in retinogeniculate refinement. *Journal of Neuroscience* **35**, 3652–3662.
- DUBIN, M.W. & CLELAND, B.G. (1977). Organization of visual inputs to interneurons of lateral geniculate nucleus of the cat. *Journal of Neurophysiology* **40**, 410–427.
- ECKER, J.L., DUMITRESCU, O.N., WONG, K.Y., ALAM, N.M., CHEN, S-K., LEGATES, T., RENNA, J.M., PRUSKY, G.T., BERSON, D.M. & HATTAR, S. (2010). Melanopsin-expressing retinal ganglion-cell photoreceptors: Cellular diversity and role in pattern vision. *Neuron* **67**, 49–60.
- ELLIS, E.M., GAUVAIN, G., SIYYER, B. & MURPHY, G.J. (2016). Shared and distinct retinal input to the mouse superior colliculus and dorsal lateral geniculate nucleus. *Journal of Neurophysiology* **116**, 602–610.
- EMERSON, V., CHALUPA, L., THOMPSON, I. & TALBOT, R. (1982). Behavioural, physiological, and anatomical consequences of monocular deprivation in the golden hamster (*Mesocricetus auratus*). *Experimental Brain Research* **45**, 168–178.
- ENGELUND, A., FAHRENKRUG, J., HARRISON, A. & HANNIBAL, J. (2010). Vesicular glutamate transporter 2 (VGLUT2) is co-stored with PACAP in projections from the rat melanopsin-containing retinal ganglion cells. *Cell and Tissue Research* **340**, 243–255.
- ERZURUMLU, R.S., JHAVERI, S. & SCHNEIDER, G.E. (1988). Distribution of morphologically different retinal axon terminals in the hamster dorsal lateral geniculate nucleus. *Brain Research* **461**, 175–181.
- ESTEVEZ, M.E., FOGERSON, P.M., ILARDI, M.C., BORGHUIS, B.G., CHAN, E., WENG, S., AUFEKORTE, O.N., DEMB, J.B. & BERSON, D.M. (2012). Form and function of the M4 cell, an intrinsically photosensitive retinal ganglion cell type contributing to geniculocortical vision. *Journal of Neuroscience* **32**, 13608–13620.
- FELDHEIM, D.A., VANDERHAEGHEN, P., HANSEN, M.J., FRISÉN, J., LU, Q., BARBACID, M. & FLANAGAN, J.G. (1998). Topographic guidance labels in a sensory projection to the forebrain. *Neuron* **21**, 1303–1313.
- FOX, M.A. & GUIDO, W. (2011). Shedding light on class-specific wiring: Development of intrinsically photosensitive retinal ganglion cell circuitry. *Molecular Neurobiology* **44**, 321–329.
- FREMEAUX, R.T., TROYER, M.D., PAHNER, I., NYGAARD, G.O., TRAN, C.H., REIMER, R.J., BELLOCCHIO, E.E., FORTIN, D., STORM-MATHISEN, J. & EDWARDS, R.H. (2001). The expression of vesicular glutamate transporters defines two classes of excitatory synapse. *Neuron* **31**, 247–260.
- FRIEDLANDER, M., LIN, C. & SHERMAN, S. (1980). Dendritic and axonal morphology of physiological classes of geniculo-cortical relay cells. In *Experimental Brain Research* (Vol. 41, No. 1, pp. A3–A3). 175 Springer Verlag, New York, US.
- FRIEDLANDER, M., LIN, C., STANFORD, L. & SHERMAN, S.M. (1981). Morphology of functionally identified neurons in lateral geniculate nucleus of the cat. *Journal of Neurophysiology* **46**, 80–129.
- FUJIYAMA, F., HIOKI, H., TOMIOKA, R., TAKI, K., TAMAMAKI, N., NOMURA, S., OKAMOTO, K. & KANEKO, T. (2003). Changes of immunocytochemical localization of vesicular glutamate transporters in the rat visual system after the retinofugal denervation. *Journal of Comparative Neurology* **465**, 234–249.
- GABBOTT, P. & BACON, S. (1994a). An oriented framework of neuronal processes in the ventral lateral geniculate nucleus of the rat demonstrated by NADPH diaphorase histochemistry and GABA immunocytochemistry. *Neuroscience* **60**, 417–440.
- GABBOTT, P.L. & BACON, S.J. (1994b). Two types of interneuron in the dorsal lateral geniculate nucleus of the rat: A combined NADPH diaphorase histochemical and GABA immunocytochemical study. *Journal of Comparative Neurology* **350**, 281–301.
- GAILLARD, F., KARTEN, H.J. & SAUVÉ, Y. (2013). Retinorecipient areas in the diurnal murine rodent *Arvicantis niloticus*: A disproportionately large superior colliculus. *Journal of Comparative Neurology* **521**, 1699–1726.
- GLEES, P. & LE GROS CLARK, W. (1941). The termination of optic fibres in the lateral geniculate body of the monkey. *Journal of Anatomy* **75**, 295.
- GODEMENT, P., SALAÜN, J. & IMBERT, M. (1984). Prenatal and postnatal development of retinogeniculate and retinocollicular projections in the mouse. *Journal of Comparative Neurology* **230**, 552–575.
- GOLDING, B., POUCHÉLON, G., BELLONE, C., MURTHY, S., DI NARDO, A.A., GOVINDAN, S., OGAWA, M., SHIMOGORI, T., LÜSCHER, C. & DAYER, A. (2014). Retinal input directs the recruitment of inhibitory interneurons into thalamic visual circuits. *Neuron* **81**, 1057–1069.
- GRANT, E., HOERDER-SUABEDISSEN, A. & MOLNÁR, Z. (2016). The regulation of corticofugal fiber targeting by retinal inputs. *Cerebral Cortex* **26**, 1336–1348.
- GRUBB, M.S. & THOMPSON, I.D. (2004). Biochemical and anatomical subdivision of the dorsal lateral geniculate nucleus in normal mice and in mice lacking the  $\beta 2$  subunit of the nicotinic acetylcholine receptor. *Vision Research* **44**, 3365–3376.
- GUIDO, W. (2008). Refinement of the retinogeniculate pathway. *The Journal of Physiology* **586**, 4357–4362.
- GUILLERY, R. (1969). The organization of synaptic interconnections in the laminae of the dorsal lateral geniculate nucleus of the cat. *Zeitschrift für Zellforschung und mikroskopische Anatomie* **96**, 1–38.
- GUILLERY, R. & HARTING, J.K. (2003). Structure and connections of the thalamic reticular nucleus: Advancing views over half a century. *Journal of Comparative Neurology* **463**, 360–371.
- HALE, P., SEFTON, A.J., BAUR, L. & COTTEE, L. (1982). Interrelations of the rat's thalamic reticular and dorsal lateral geniculate nuclei. *Experimental Brain Research* **45**, 217–229.
- HALLANGER, A.E., LEVEY, A.I., LEE, H.J., RYE, D.B. & WAINER, B.H. (1987). The origins of cholinergic and other subcortical afferents to the thalamus in the rat. *Journal of Comparative Neurology* **262**, 105–124.
- HAMMER, S., CARRILLO, G.L., GOVINDAIAH, G., MONAVARFESHANI, A., BIRCHER, J.S., SU, J., GUIDO, W. & FOX, M.A. (2014). Nuclei-specific differences in nerve terminal distribution, morphology, and development in mouse visual thalamus. *Neural Development* **9**, 1.
- HAMMER, S., MONAVARFESHANI, A., LEMON, T., SU, J. & FOX, M.A. (2015). Multiple retinal axons converge onto relay cells in the adult mouse thalamus. *Cell Reports* **12**, 1575–1583.
- HARRINGTON, M., DECOURSEY, P., BRUCE, D. & BUGGY, J. (1987). Circadian pacemaker (SCN) transplants into lateral ventricles fail to restore locomotor rhythmicity in arrhythmic hamsters. *Soc Neurosci Abstr* **13**, 465–472.
- HARRINGTON, M.E. (1997). The ventral lateral geniculate nucleus and the intergeniculate leaflet: Interrelated structures in the visual and circadian systems. *Neuroscience & Biobehavioral Reviews* **21**, 705–727.
- HARRINGTON, M.E. & RUSAK, B. (1986). Lesions of the thalamic intergeniculate leaflet alter hamster circadian rhythms. *Journal of Biological Rhythms* **1**, 309–325.

- HARTING, J.K., HUERTA, M.F., HASHIKAWA, T. & VAN LIESHOUT, D.P. (1991a). Projection of the mammalian superior colliculus upon the dorsal lateral geniculate nucleus: Organization of tectogeniculate pathways in nineteen species. *Journal of Comparative Neurology* **304**, 275–306.
- HARTING, J.K., VAN LIESHOUT, D., HASHIKAWA, T. & WEBER, J. (1991b). The paravigenogulate projection: Connectional studies in eight mammals. *Journal of Comparative Neurology* **305**, 559–581.
- HATTAR, S., KUMAR, M., PARK, A., TONG, P., TUNG, J., YAU, K.W. & BERSON, D.M. (2006). Central projections of melanopsin-expressing retinal ganglion cells in the mouse. *Journal of Comparative Neurology* **497**, 326–349.
- HICKEY, T. & SPEAR, P. (1976). Retinogeniculate projections in hooded and albino rats: An autoradiographic study. *Experimental Brain Research* **24**, 523–529.
- HOLCOMBE, V. & GUILLERY, R. (1984). The organization of retinal maps within the dorsal and ventral lateral geniculate nuclei of the rabbit. *Journal of Comparative Neurology* **225**, 469–491.
- HONG, Y.K. & CHEN, C. (2011). Wiring and rewiring of the retinogeniculate synapse. *Current Opinion in Neurobiology* **21**, 228–237.
- HOREL, J.A. (1968). Effects of subcortical lesions on brightness discrimination acquired by rats without visual cortex. *Journal of Comparative and Physiological Psychology* **65**, 103.
- HOROWITZ, S.S., BLANCHARD, J.H. & MORIN, L.P. (2004). Intergeniculate leaflet and ventral lateral geniculate nucleus afferent connections: An anatomical substrate for functional input from the vestibulo-visuomotor system. *Journal of Comparative Neurology* **474**, 227–245.
- HUBERMAN, A.D., FELLER, M.B. & CHAPMAN, B. (2008a). Mechanisms underlying development of visual maps and receptive fields. *Annual Review of Neuroscience* **31**, 479–509.
- HUBERMAN, A.D., MANU, M., KOCH, S.M., SUSMAN, M.W., LUTZ, A.B., ULLIAN, E.M., BACCUS, S.A. & BARRES, B.A. (2008b). Architecture and activity-mediated refinement of axonal projections from a mosaic of genetically identified retinal ganglion cells. *Neuron* **59**, 425–438.
- HUBERMAN, A.D., WEI, W., ELSTROTT, J., STAFFORD, B.K., FELLER, M.B. & BARRES, B.A. (2009). Genetic identification of an on-off direction-selective retinal ganglion cell subtype reveals a layer-specific corticofugal map of posterior motion. *Neuron* **62**, 327–334.
- INAMURA, N., ONO, K., TAKEBAYASHI, H., ZALC, B. & IKENAKA, K. (2011). Olig2 lineage cells generate GABAergic neurons in the prethalamic nuclei, including the zona incerta, ventral lateral geniculate nucleus and reticular thalamic nucleus. *Developmental Neuroscience* **33**, 118–129.
- IRVIN, G.E., CASAGRANDE, V.A. & NORTON, T.T. (1993). Center/surround relationships of magnocellular, parvocellular, and koniocellular relay cells in primate lateral geniculate nucleus. *Visual Neuroscience* **10**, 363–373.
- JACOBS, E.C., CAMPAGNONI, C., KAMPE, K., REYES, S.D., KALRA, V., HANDLEY, V., XIE, Y.Y., HONG-HU, Y., SPREUR, V. & FISHER, R.S. (2007). Visualization of corticofugal projections during early cortical development in a  $\tau$ -GFP-transgenic mouse. *European Journal of Neuroscience* **25**, 17–30.
- JAGER, P., YE, Z., YU, X., ZAGORAIOU, L., PREKOP, H-T., PARTANEN, J., JESSELL, T.M., WISDEN, W., BRICKLEY, S.G. & DELOGU, A. (2016). Tectal-derived interneurons contribute to phasic and tonic inhibition in the visual thalamus. *Nature Communications* **7**, 13579.
- JAUBERT-MIAZZA, L., GREEN, E., LO, F., BUI, K., MILLS, J. & GUIDO, W. (2005). Structural and functional composition of the developing retinogeniculate pathway in the mouse. *Visual Neuroscience* **22**, 661.
- JONES, E.G. (2012). *The Thalamus*. Springer Science & Business Media, Berlin, Germany.
- JONES, E.G. & RUBENSTEIN, J.L. (2004). Expression of regulatory genes during differentiation of thalamic nuclei in mouse and monkey. *Journal of Comparative Neurology* **477**, 55–80.
- JURGENS, C.W., BELL, K.A., MCQUISTON, A.R. & GUIDO, W. (2012). Optogenetic stimulation of the corticothalamic pathway affects relay cells and GABAergic neurons differently in the mouse visual thalamus. *PLoS One* **7**, e45717.
- KAAS, J., GUILLERY, R. & ALLMAN, J. (1972). Some principles of organization in the dorsal lateral geniculate nucleus. *Brain, Behavior and Evolution* **6**, 283–299.
- KAY, J.N., DE LA HUERTA, I., KIM, I-J., ZHANG, Y., YAMAGATA, M., CHU, M.W., MEISTER, M. & SANES, J.R. (2011). Retinal ganglion cells with distinct directional preferences differ in molecular identity, structure, and central projections. *The Journal of Neuroscience* **31**, 7753–7762.
- KIM, I-J., ZHANG, Y., MEISTER, M. & SANES, J.R. (2010). Laminar restriction of retinal ganglion cell dendrites and axons: Subtype-specific developmental patterns revealed with transgenic markers. *The Journal of Neuroscience* **30**, 1452–1462.
- KIM, I-J., ZHANG, Y., YAMAGATA, M., MEISTER, M. & SANES, J.R. (2008). Molecular identification of a retinal cell type that responds to upward motion. *Nature* **452**, 478–482.
- KOLMAC, C.I., POWER, B.D. & MITROFANIS, J. (2000). Dorsal thalamic connections of the ventral lateral geniculate nucleus of rats. *Journal of Neurocytology* **29**, 31–41.
- KOPP, M.D., MEISSL, H., DEGHANI, F. & KORF, H.W. (2001). The pituitary adenylate cyclase-activating polypeptide modulates glutamatergic calcium signalling: Investigations on rat suprachiasmatic nucleus neurons. *Journal of Neurochemistry* **79**, 161–171.
- KRAHE, T.E., EL-DANAF, R.N., DILGER, E.K., HENDERSON, S.C. & GUIDO, W. (2011). Morphologically distinct classes of relay cells exhibit regional preferences in the dorsal lateral geniculate nucleus of the mouse. *The Journal of Neuroscience* **31**, 17437–17448.
- LEGG, C. & COWEY, A. (1977a). Effects of subcortical lesions on visual intensity discriminations in rats. *Physiology & Behavior* **19**, 635–646.
- LEGG, C. & COWEY, A. (1977b). The role of the ventral lateral geniculate nucleus and posterior thalamus in intensity discrimination in rats. *Brain Research* **123**, 261–273.
- LEIST, M., DATUNASHVILLI, M., KANYSHKOVA, T., ZOBEIRI, M., AISSAOUI, A., CERINA, M., ROMANELLI, M.N., PAPE, H-C. & BUDDÉ, T. (2016). Two types of interneurons in the mouse lateral geniculate nucleus are characterized by different h-current density. *Scientific Reports* **6**, 24904.
- LEWANDOWSKI, M.H. & USAREK, A. (2002). Effects of intergeniculate leaflet lesions on circadian rhythms in the mouse. *Behavioural Brain Research* **128**, 13–17.
- LING, C., HENDRICKSON, M.L. & KALIL, R.E. (2012). Morphology, classification, and distribution of the projection neurons in the dorsal lateral geniculate nucleus of the rat. *PLoS One* **7**, e49161.
- LÓPEZ-BENDITO, G. & MOLNÁR, Z. (2003). Thalamocortical development: How are we going to get there? *Nature Reviews Neuroscience* **4**, 276–289.
- LUND, R. & CUNNINGHAM, T. (1972). Aspects of synaptic and laminar organization of the mammalian lateral geniculate body. *Investigative Ophthalmology* **11**, 291–302.
- MACKAY-SIM, A., SEFTON, A.J. & MARTIN, P.R. (1983). Subcortical projections to lateral geniculate and thalamic reticular nuclei in the hooded rat. *Journal of Comparative Neurology* **213**, 24–35.
- MARTERSTECK, E.M., HIROKAWA, K.E., EVARTS, M., BERNARD, A., DUAN, X., LI, Y., NG, L., OH, S.W., OUELLETTE, B. & ROYALL, J.J. (2017). Diverse central projection patterns of retinal ganglion cells. *Cell Reports* **18**, 2058–2072.
- MARTINEZ, L.M., MOLANO-MAZÓN, M., WANG, X., SOMMER, F.T. & HIRSCH, J.A. (2014). Statistical wiring of thalamic receptive fields optimizes spatial sampling of the retinal image. *Neuron* **81**, 943–956.
- MATUTE, C. & STREIT, P. (1985). Selective retrograde labeling with D-[3H]-aspartate in afferents to the mammalian superior colliculus. *Journal of Comparative Neurology* **241**, 34–49.
- MCCORMICK, D.A. (1992). Neurotransmitter actions in the thalamus and cerebral cortex and their role in neuromodulation of thalamocortical activity. *Progress in Neurobiology* **39**, 337–388.
- MICHEL, S., ITRI, J., HAN, J.H., GNITCZYNSKI, K. & COLWELL, C.S. (2006). Regulation of glutamatergic signalling by PACAP in the mammalian suprachiasmatic nucleus. *BMC Neuroscience* **7**, 15.
- MIKKELSEN, J. (1992). The organization of the crossed geniculogeniculate pathway of the rat: A Phaseolus vulgaris-leucoagglutinin study. *Neuroscience* **48**, 953–962.
- MIZE, R.R. & HORNER, L.H. (1984). Retinal synapses of the cat medial interlaminar nucleus and ventral lateral geniculate nucleus differ in size and synaptic organization. *Journal of Comparative Neurology* **224**, 579–590.
- MONTERA, V. & ZEMPEL, J. (1985). Evidence for two types of GABA-containing interneurons in the  $\alpha$ -laminae of the cat lateral geniculate nucleus: A double-label HRP and GABA-immunocytochemical study. *Experimental Brain Research* **60**, 603–609.
- MONTERO, V. & SINGER, W. (1985). Ultrastructural identification of somata and neural processes immunoreactive to antibodies against glutamic acid decarboxylase (GAD) in the dorsal lateral geniculate nucleus of the cat. *Experimental Brain Research* **59**, 151–165.
- MOORE, R.Y. & CARD, J.P. (1994). Intergeniculate leaflet: An anatomically and functionally distinct subdivision of the lateral geniculate complex. *Journal of Comparative Neurology* **344**, 403–430.

- MOORE, R.Y. & SPEH, J.C. (1993). GABA is the principal neurotransmitter of the circadian system. *Neuroscience Letters* **150**, 112–116.
- MOORE, R.Y., WEIS, R. & MOGA, M.M. (2000). Efferent projections of the intergeniculate leaflet and the ventral lateral geniculate nucleus in the rat. *Journal of Comparative Neurology* **420**, 398–418.
- MORGAN, J.L., BERGER, D.R., WETZEL, A.W. & LICHTMAN, J.W. (2016). The fuzzy logic of network connectivity in mouse visual thalamus. *Cell* **165**, 192–206.
- MORIN, L. & BLANCHARD, J. (1998). Interconnections among nuclei of the subcortical visual shell: The intergeniculate leaflet is a major constituent of the hamster subcortical visual system. *Journal of Comparative Neurology* **396**, 288–309.
- MORIN, L. & BLANCHARD, J. (1999). Forebrain connections of the hamster intergeniculate leaflet: Comparison with those of ventral lateral geniculate nucleus and retina. *Visual Neuroscience* **16**, 1037–1054.
- MORIN, L.P. (2013). Neuroanatomy of the extended circadian rhythm system. *Experimental Neurology* **243**, 4–20.
- MORIN, L.P. & STUDHOLME, K.M. (2014). Retinofugal projections in the mouse. *Journal of Comparative Neurology* **522**, 3733–3753.
- MUIR-ROBINSON, G., HWANG, B.J. & FELLER, M.B. (2002). Retinogeniculate axons undergo eye-specific segregation in the absence of eye-specific layers. *The Journal of Neuroscience* **22**, 5259–5264.
- NAKAGAWA, Y. & SHIMOGORI, T. (2012). Diversity of thalamic progenitor cells and postmitotic neurons. *European Journal of Neuroscience* **35**, 1554–1562.
- NAKAMURA, H. & KAWAMURA, S. (1988). The ventral lateral geniculate nucleus in the cat: Thalamic and commissural connections revealed by the use of WGA-HRP transport. *Journal of Comparative Neurology* **277**, 509–528.
- NASSI, J.J. & CALLAWAY, E.M. (2009). Parallel processing strategies of the primate visual system. *Nature Reviews Neuroscience* **10**, 360–372.
- NIIMI, K., KANASEKI, T. & TAKIMOTO, T. (1963). The comparative anatomy of the ventral nucleus of the lateral geniculate body in mammals. *Journal of Comparative Neurology* **121**, 313–323.
- OLSEN, S.R., BORTONE, D.S., ADESNIK, H. & SCANZIANI, M. (2012). Gain control by layer six in cortical circuits of vision. *Nature* **483**, 47–52.
- OSTERHOUT, J.A., JOSTEN, N., YAMADA, J., PAN, F., WU, S.-W., NGUYEN, P.L., PANAGIOTAKOS, G., INOUE, Y.U., EGUSA, S.F. & VOLGYI, B. (2011). Cadherin-6 mediates axon-target matching in a non-image-forming visual circuit. *Neuron* **71**, 632–639.
- PAPADOPOULOS, G.C. & PARNAVELAS, J.G. (1990a). Distribution and synaptic organization of dopaminergic axons in the lateral geniculate nucleus of the rat. *Journal of Comparative Neurology* **294**, 356–361.
- PAPADOPOULOS, G.C. & PARNAVELAS, J.G. (1990b). Distribution and synaptic organization of serotonergic and noradrenergic axons in the lateral geniculate nucleus of the rat. *Journal of Comparative Neurology* **294**, 345–355.
- PETROF, I. & SHERMAN, S.M. (2013). Functional significance of synaptic terminal size in glutamatergic sensory pathways in thalamus and cortex. *The Journal of physiology* **591**, 3125–3131.
- PFEIFFENBERGER, C., YAMADA, J. & FELDHEIM, D.A. (2006). Ephrin-As and patterned retinal activity act together in the development of topographic maps in the primary visual system. *The Journal of Neuroscience* **26**, 12873–12884.
- PINAULT, D. (2004). The thalamic reticular nucleus: Structure, function and concept. *Brain Research Reviews* **46**, 1–31.
- PISCOPO, D.M., EL-DANAF, R.N., HUBERMAN, A.D. & NIELL, C.M. (2013). Diverse visual features encoded in mouse lateral geniculate nucleus. *The Journal of Neuroscience* **33**, 4642–4656.
- PUELLES, L. & RUBENSTEIN, J.L. (2003). Forebrain gene expression domains and the evolving prosomeric model. *Trends in Neurosciences* **26**, 469–476.
- RAFOLS, J.A. & VALVERDE, F. (1973). The structure of the dorsal lateral geniculate nucleus in the mouse. A golgi and electron microscopic study. *Journal of Comparative Neurology* **150**, 303–331.
- REBSAM, A., BHANSALI, P. & MASON, C.A. (2012). Eye-specific projections of retinogeniculate axons are altered in albino mice. *Journal of Neuroscience* **32**, 4821–4826.
- REBSAM, A., PETROS, T.J. & MASON, C.A. (2009). Switching retinogeniculate axon laterality leads to normal targeting but abnormal eye-specific segregation that is activity dependent. *Journal of Neuroscience* **29**, 14855–14863.
- REESE, B. (1988). 'Hidden lamination' in the dorsal lateral geniculate nucleus: The functional organization of this thalamic region in the rat. *Brain Research Reviews* **13**, 119–137.
- REESE, B. & COWEY, A. (1983). Projection lines and the ipsilateral retinogeniculate pathway in the hooded rat. *Neuroscience* **10**, 1233–1247.
- RIBAK, C.E. & PETERS, A. (1975). An autoradiographic study of the projections from the lateral geniculate body of the rat. *Brain Research* **92**, 341–368.
- RIVLIN-ETZION, M., ZHOU, K., WEI, W., ELSTROTT, J., NGUYEN, P.L., BARRÉS, B.A., HUBERMAN, A.D. & FELLER, M.B. (2011). Transgenic mice reveal unexpected diversity of on-off direction-selective retinal ganglion cell subtypes and brain structures involved in motion processing. *The Journal of Neuroscience* **31**, 8760–8769.
- ROMPANI, S.B., MÜLLNER, F.E., WANNER, A., ZHANG, C., ROTH, C.N., YONEHARA, K. & ROSKA, B. (2017). Different modes of visual integration in the lateral geniculate nucleus revealed by single-cell-initiated transsynaptic tracing. *Neuron* **93**, 767–776. e766.
- ROTH, M.M., DAHMEN, J.C., MUIR, D.R., IMHOF, F., MARTINI, F.J. & HOFER, S.B. (2016). Thalamic nuclei convey diverse contextual information to layer 1 of visual cortex. *Nature Neuroscience* **19**, 299–307.
- SANES, J.R. & MASLAND, R.H. (2015). The types of retinal ganglion cells: Current status and implications for neuronal classification. *Annual Review of Neuroscience* **38**, 221–246.
- SCHMIDT, T.M., CHEN, S.-K. & HATTAR, S. (2011). Intrinsically photosensitive retinal ganglion cells: Many subtypes, diverse functions. *Trends in Neurosciences* **34**, 572–580.
- SEABROOK, T.A., EL-DANAF, R.N., KRAHE, T.E., FOX, M.A. & GUIDO, W. (2013a). Retinal input regulates the timing of corticogeniculate innervation. *The Journal of Neuroscience* **33**, 10085–10097.
- SEABROOK, T.A., KRAHE, T.E., GOVINDAIAH, G. & GUIDO, W. (2013b). Interneurons in the mouse visual thalamus maintain a high degree of retinal convergence throughout postnatal development. *Neural Development* **8**, 1.
- SHANKS, J.A., ITO, S., SCHAEVITZ, L., YAMADA, J., CHEN, B., LITKE, A. & FELDHEIM, D.A. (2016). Corticothalamic axons are essential for retinal ganglion cell axon targeting to the mouse dorsal lateral geniculate nucleus. *The Journal of Neuroscience* **36**, 5252–5263.
- SHERMAN, S.M. (1985). Functional organization of the W-, X-, and Y-cell pathways in the cat: A review and hypothesis. *Progress in Psychobiology and Physiological Psychology* **11**, 233–314.
- SHERMAN, S.M. (2004). Interneurons and triadic circuitry of the thalamus. *Trends in Neurosciences* **27**, 670–675.
- SHERMAN, S.M. (2005). Thalamic relays and cortical functioning. *Progress in Brain Research* **149**, 107–126.
- SHERMAN, S.M. (2016). Thalamus plays a central role in ongoing cortical functioning. *Nature Neuroscience* **16**, 533–541.
- SHERMAN, S.M. & GULLERY, R. (2002). The role of the thalamus in the flow of information to the cortex. *Philosophical Transactions of the Royal Society of London Series B: Biological Sciences* **357**, 1695–1708.
- SINGH, R., SU, J., BROOKS, J., TERAUCHI, A., UMEMORI, H. & FOX, M.A. (2012). Fibroblast growth factor 22 contributes to the development of retinal nerve terminals in the dorsal lateral geniculate nucleus. *Frontiers in molecular neuroscience* **4**, 61.
- STELZNER, D.J., BAISDEN, R.H. & GOODMAN, D.C. (1976). The ventral lateral geniculate nucleus, pars lateralis of the rat. *Cell and Tissue Research* **170**, 435–454.
- STEVENS, B., ALLEN, N.J., VAZQUEZ, L.E., HOWELL, G.R., CHRISTOPHERSON, K.S., NOURI, N., MICHEVA, K.D., MEHALOW, A.K., HUBERMAN, A.D. & STAFFORD, B. (2007). The classical complement cascade mediates CNS synapse elimination. *Cell* **131**, 1164–1178.
- SU, J., HANER, C.V., IMBERY, T.E., BROOKS, J.M., MORHARDT, D.R., GORSE, K., GUIDO, W. & FOX, M.A. (2011). Reelin is required for class-specific retinogeniculate targeting. *The Journal of Neuroscience* **31**, 575–586.
- SU, J., KLEMM, M.A., JOSEPHSON, A.M. & FOX, M.A. (2013). Contributions of VLDLR and LRP8 in the establishment of retinogeniculate projections. *Neural Development* **8**, 1.
- SWANSON, L., COWAN, W. & JONES, E. (1974). An autoradiographic study of the efferent connections of the ventral lateral geniculate nucleus in the albino rat and the cat. *Journal of Comparative Neurology* **156**, 143–163.
- TAYLOR, A., JEFFERY, G. & LIEBERMAN, A. (1986). Subcortical afferent and efferent connections of the superior colliculus in the rat and comparisons between albino and pigmented strains. *Experimental Brain Research* **62**, 131–142.
- THANKACHAN, S. & RUSAK, B. (2005). Juxtacellular recording/labeling analysis of physiological and anatomical characteristics of rat intergeniculate leaflet neurons. *The Journal of Neuroscience* **25**, 9195–9204.

- THOMPSON, A.D., PICARD, N., MIN, L., FAGIOLINI, M. & CHEN, C. (2016). Cortical feedback regulates feedforward retinogeniculate refinement. *Neuron* **91**, 1021–1033.
- TOWNS, L.C., BURTON, S., KIMBERLY, C. & FETTERMAN, M. (1982). Projections of the dorsal lateral geniculate and lateral posterior nuclei to visual cortex in the rabbit. *Journal of Comparative Neurology* **210**, 87–98.
- TRIPLETT, J.W., WEI, W., GONZALEZ, C., SWEENEY, N.T., HUBERMAN, A.D., FELLER, M.B. & FELDHEIM, D.A. (2014). Dendritic and axonal targeting patterns of a genetically-specified class of retinal ganglion cells that participate in image-forming circuits. *Neural Development* **9**, 2.
- VAN HOOSER, S.D. & NELSON, S.B. (2006). The squirrel as a rodent model of the human visual system. *Visual Neuroscience* **23**, 765–778.
- VIROLAINEN, S-M., ACHIM, K., PELTOPURO, P., SALMINEN, M. & PARTANEN, J. (2012). Transcriptional regulatory mechanisms underlying the GABAergic neuron fate in different diencephalic prosomeres. *Development* **139**, 3795–3805.
- VRANG, N., MROSOVSKY, N. & MIKKELSEN, J.D. (2003). Afferent projections to the hamster intergeniculate leaflet demonstrated by retrograde and anterograde tracing. *Brain Research Bulletin* **59**, 267–288.
- VUE, T.Y., AAKER, J., TANIGUCHI, A., KAZEMZADEH, C., SKIDMORE, J.M., MARTIN, D.M., MARTIN, J.F., TREIER, M. & NAKAGAWA, Y. (2007). Characterization of progenitor domains in the developing mouse thalamus. *Journal of Comparative Neurology* **505**, 73–91.
- WANG, S., EISENBACK, M., DATSKOVSKAIA, A., BOYCE, M. & BICKFORD, M.E. (2002). GABAergic pretectal terminals contact GABAergic interneurons in the cat dorsal lateral geniculate nucleus. *Neuroscience Letters* **323**, 141–145.
- WEYAND, T.G. (2016). The multifunctional lateral geniculate nucleus. *Reviews in the neurosciences* **27**, 135–157.
- WILSON, J., FRIEDLANDER, M. & SHERMAN, S. (1984). Fine structural morphology of identified X- and Y-cells in the cat's lateral geniculate nucleus. *Proceedings of the Royal Society of London Series B, Biological Sciences* **221**, 411–436.
- XU, H-P., FURMAN, M., MINEUR, Y.S., CHEN, H., KING, S.L., ZENISEK, D., ZHOU, Z.J., BUTTS, D.A., TIAN, N., PICCIOTTO, M.R. & CRAIR, M.C. (2011). An instructive role for patterned spontaneous retinal activity in mouse visual map development. *Neuron* **70**, 1115–1127.
- YUGE, K., KATAOKA, A., YOSHIDA, A.C., ITOH, D., AGGARWAL, M., MORI, S., BLACKSHAW, S. & SHIMOGORI, T. (2011). Region-specific gene expression in early postnatal mouse thalamus. *Journal of Comparative Neurology* **519**, 544–561.
- ZHANG, J., ACKMAN, J.B., XU, H-P. & CRAIR, M.C. (2012). Visual map development depends on the temporal pattern of binocular activity in mice. *Nature Neuroscience* **15**, 298–307.